

Moduli of rational fibrations

Edoardo Sernesi*

Abstract

We prove results implying, under mild restrictions, the upper bound $\gamma \leq 19$ for the geometric genus of a nodal irreducible curve whose normalization has general moduli and varying in a non-trivial linear system of curves of the same geometric genus on some non-ruled surface. This gives some evidence for the non-uniruledness of $\overline{M}_{20}$ and $\overline{M}_{21}$. The proofs use deformation theory of surfaces fibered over the projective line.

Introduction

It is known that $\overline{M}_g$, the coarse moduli space of stable curves of genus g , has Kodaira dimension ≥ 2 if $g = 23$ ([12]), and is of general type for all other values of $g \geq 22$ ([16], [11], [13]). In particular $\overline{M}_g$ is not uniruled when $g \geq 22$. This last property implies the following:

Proposition 0.1 *If $g \geq 22$, then a general curve of genus g does not occur in any non-trivial linear system on any non-ruled surface.*

(see the Introduction to [16] for an outline of the proof). Since we don't yet know all the values of g for which $\overline{M}_g$ is not uniruled, it is of some interest to look for a proof of Proposition 0.1, and for a generalization of it to smaller values of g , which does not make use of any information about the Kodaira dimension of $\overline{M}_g$. The present work is a step in this direction.

The first most natural attempt to obtain a general curve of a given genus g occurring in a non-trivial linear system on a surface, would be to look for

*Member of GNSAGA-INDAM.

a surface Y containing a linear system Λ of curves of genus g such that the family of curves parametrized by Λ defines a dominant rational map to $\overline{\mathcal{M}}_g$. The existence of such a configuration would imply the unirationality of $\overline{\mathcal{M}}_g$, and not only its uniruledness. This point of view goes back to Severi and B. Segre, who studied rational and unirational families of plane curves of given geometric genus and having general moduli (see [4], [6] and references therein). Here we are interested in something weaker, since we want to allow the surface Y to vary in a family carrying along the linear system.

More precisely, to say that “a general curve of genus g occurs in a non-trivial linear system on some surface”, means that we can find a family of pairs (C_t, Y_t) , where t varies in some irreducible parameter variety M , and where C_t is a nonsingular projective connected curve of genus g , Y_t is a nonsingular projective surface, so that

$$C_t \subset Y_t,$$

$$\dim(|C_t|) \geq 1 \text{ for all } t \in M,$$

$$[C_t] \text{ spans an open subset of } \overline{\mathcal{M}}_g \text{ as } t \text{ varies in } M.$$

To a general linear pencil contained in any of the linear systems $|C_t|$ one associates a fibration

$$f : X \rightarrow \mathbb{P}^1$$

in curves of genus g , so that, conversely, the question of existence of the family (C_t, Y_t) translates into a question about deformations of a fibration f .

Given a fibration f , we study its local deformation theory in the sense of Horikawa. We do it more generally when the base is a nonsingular projective connected curve S , not necessarily $\mathbb{P}^1$, and we obtain some estimates on the number of moduli of f .

In the case $S = \mathbb{P}^1$ (a *rational fibration*) we introduce the notion of *free rational fibration*, and we prove that those rational fibrations which can be deformed so as to contain a general curve of genus g as a fibre, are free (Theorem 2.6). This notion is analogous to the one of “free rational curve” in a nonsingular variety M [3]; in fact, if it is semistable then a free fibration defines a free rational curve in the moduli stack $\overline{\mathcal{M}}_g$ of stable curves of genus g . All the above leads to necessary numerical conditions for a surface Y to carry a linear pencil whose general element is a nonsingular curve of genus g , and defining a free fibration (Theorem 3.2). The most notable of such

estimates is the following one:

$$10\chi(\mathcal{O}_Y) - 2K_Y^2 \geq 4(g-1) - C^2$$

where C is a curve of the pencil, which holds provided $|K_Y - C| = \emptyset$. Note that the left hand side equals $-\chi(T_Y)$, and therefore it coincides with $\mu(Y) := h^1(T_Y) - h^2(T_Y)$, the expected number of moduli of Y , in case $H^0(T_Y) = 0$. Therefore, in such a case, we get an explicit lower bound on $\mu(Y)$ as a necessary condition for the surface to carry a general nonsingular curve of genus g moving in a non-trivial linear system. Conversely, in order to bound the genus from above, we need that $-\chi(T_Y)$ does not get too big or, what is the same, that K_Y^2 is as large as possible. This suggests to develop our analysis on a *minimal* surface where, if we want to maintain our analysis completely general, we have to consider *curves with arbitrary singularities* having general normalization and moving in a non-trivial linear system of curves of the same geometric genus. This requires an appropriate generalization of the above inequality to linear pencils of singular curves. This is done in §3, Proposition 3.5, under the restriction that the curves have only nodes (ordinary double points) as singularities. Using these inequalities we are able to give upper bounds on the genus of a nodal curve whose normalization has general moduli, varying in a non-trivial linear system on some non-ruled surface Y .

The following preliminary results cover all the cases of curves on surfaces of low Kodaira dimension:

$Y = \mathbb{P}^2$. Then every linear pencil of nodal curves whose normalization has general moduli consists of curves of degree ≤ 8 and geometric genus ≤ 9 (Theorem 4.1). This result is essentially classical and does not need the deformation theory of fibrations for its proof.

$Y = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(n))$, $n \geq 0$, $n \neq 1$, is a minimal rational surface. Then every linear pencil of nodal curves whose normalization has general moduli consists of curves of geometric genus ≤ 14 (Theorem 4.2).

Y is a minimal non-ruled and non-rational surface of geometric genus $p_g \leq 2$. Then every linear pencil of nodal curves whose normalization has general moduli consists of curves of geometric genus

$$\gamma \leq \begin{cases} 6 & \text{if } p_g = 0 \\ 11 & \text{if } p_g = 1 \\ 16 & \text{if } p_g = 2 \end{cases}$$

(Theorem 5.1).

Y is a minimal elliptic surface of nonnegative Kodaira dimension, Then every linear pencil of nodal curves whose normalization has general moduli consists of curves of geometric genus $\gamma \leq 16$ (Theorem 5.3).

Some of the above results are probably folklore but, with the exception of the cases of $\mathbb{P}^2$ and of K3 surfaces, we were unable to locate them in the literature. Our main results concern curves on surfaces of general type. The first one is the following (Theorem 6.1):

Theorem 0.2 *Let Y be a minimal surface of general type, $C \subset Y$ a nodal irreducible curve of geometric genus γ whose normalization has general moduli. Denote by $|C(-2p_1 - \dots - 2p_\delta)|$ be the linear system of curves in $|C|$ singular at the nodes $p_1, \dots, p_\delta$ of C . Then:*

- if $K_Y^2 \geq 3\chi(\mathcal{O}_Y) - 10$ and $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$ then $\gamma \leq 19$,
- if $K_Y^2 \geq 4\chi(\mathcal{O}_Y) - 16$ and $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$ then $\gamma \leq 15$,
- if $K_Y^2 \geq 4\chi(\mathcal{O}_Y) - 16$, $|K_Y - C| = \emptyset$ and $\dim(|C(-2p_1 - \dots - 2p_\delta)|) = 1$ then $\gamma \leq 19$.

As a notable special case we obtain (Corollary 6.2):

Corollary 0.3 *Let Y be a minimal surface of general type such that the canonical map is birational onto its image. Suppose that $C \subset Y$ is a nodal irreducible curve of geometric genus γ whose normalization has general moduli and such that $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$, where $p_1, \dots, p_\delta$ are the nodes of C . Then $\gamma \leq 19$.*

The remaining cases of minimal surfaces of general type with small K^2 are considered by the next result:

Theorem 0.4 *Let Y be a minimal surface of general type such that*

$$K_Y^2 < 3\chi(\mathcal{O}_Y) - 10$$

and $C \subset Y$ a a nodal irreducible curve of geometric genus γ whose normalization has general moduli, having nodes $p_1, \dots, p_\delta$, such that $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 1$. Then (under a technical assumption, see the statement of Theorem 6.5) Y is regular and $\gamma \leq 14$.

We do not claim that all these statements are sharp. In the recent paper [9] the authors construct a remarkable example in the case $\gamma = 15$ which turns out to be borderline in one of the cases of Theorem 0.2; so at least in this case our result is sharp.

Coming back to our original motivation of proving a generalization of Proposition 0.1, in relation with the non-uniruledness of $\overline{M}_g$, we can summarize our results as follows. Our analysis does not exclude the existence of non-trivial linear systems of general curves of geometric genus ≥ 20 *with arbitrary singularities*, on minimal surfaces, but only of such linear systems consisting of nodal curves. Apart from this, and from the condition that the linear system has dimension ≥ 2 in some cases, our analysis is otherwise quite general, and gives, in particular, some evidence for the non-uniruledness of $\overline{M}_{20}$ and $\overline{M}_{21}$.

The paper is divided into six sections. Section 1 contains generalities about fibrations. Section 2 is devoted to the deformation theory of fibrations, and Section 3 applies this analysis to the fibrations deduced from linear pencils of curves on surfaces, obtaining the key inequalities needed. In Section 4 we discuss curves on rational surfaces, and in Sections 5 we consider curves on surfaces of low Kodaira dimension. In Section 6 we consider the main case of curves on minimal surfaces of general type.

Acknowledgements. I am grateful to the Institut Mittag-Leffler (Djursholm, Sweden) for support and hospitality during the preparation of part of this work. I thank A. Bruno, C. Ciliberto and M. Roth for interesting conversations related with this work, and F. Catanese for helpful remarks.

1 Fibrations

We work over an algebraically closed field $\mathbf{k}$ of characteristic 0. We will refer to [17] for all notions and terminology not explicitly introduced.

If

$$\varphi : W \rightarrow U$$

is a morphism of algebraic schemes, and $u \in U$, we denote by $W(u)$ the scheme-theoretic fibre of φ over u .

A (-1) -*curve* (resp. a (-2) -*curve*) on a projective nonsingular surface Z is a nonsingular connected rational curve with self-intersection -1 (resp. -2). We will denote by K_Z a canonical divisor on Z . By a *minimal surface*

we mean, as customary, a projective nonsingular connected surface without (-1) -curves.

By a *fibration* we mean a surjective morphism

$$f : X \rightarrow S$$

with connected fibres from a projective nonsingular surface to a projective nonsingular connected curve. We will denote by

$g =$ the genus of the general fibre. *We will always assume $g \geq 2$.*

$b =$ the genus of S .

A fibration is called:

- *relatively minimal* if there are no (-1) -curves contained in any of its fibres.
- *semistable* if it is relatively minimal and every fibre has at most nodes as singularities.
- *isotrivial* if all its nonsingular fibres are mutually isomorphic; equivalently, if two general nonsingular fibres of f are mutually isomorphic (the equivalence of the two formulations follows from the separatedness of $\overline{M}_g$).

Sheaves of differentials will be denoted with the symbol Ω and dualizing sheaves with the symbol ω . Let $f : X \rightarrow S$ be any fibration. Since both X and S are nonsingular we have:

$$\wedge^2 \Omega_X^1 = \omega_X, \quad \Omega_S^1 = \omega_S$$

Moreover f is a relative complete intersection morphism and it follows that

$$\omega_{X/S} = \omega_X \otimes f^* \omega_S^{-1} \tag{1}$$

(see [18], Corollary (24)). We have an exact sequence

$$0 \rightarrow f^* \omega_S \rightarrow \Omega_X^1 \rightarrow \Omega_{X/S}^1 \rightarrow 0 \tag{2}$$

(which is exact on the left because the first homomorphism is injective on a dense open set and $f^*\omega_S$ is locally free). If we dualize the sequence (2) we obtain the exact sequence:

$$0 \rightarrow T_{X/S} \rightarrow T_X \rightarrow f^*T_S \rightarrow N \rightarrow 0 \quad (3)$$

where we have denoted

$$T_{X/S} := \text{Hom}_{\mathcal{O}_X}(\Omega_{X/S}^1, \mathcal{O}_X)$$

and

$$N := \text{Ext}_{\mathcal{O}_X}^1(\Omega_{X/S}^1, \mathcal{O}_X) \quad (4)$$

The sequence (3) shows, in particular, that N is the normal sheaf of f , and also the first relative cotangent sheaf $T_{X/S}^1$ of f (see [24]). In particular, N is supported on the set of singular points of the fibres of f .

Moreover, as remarked in [25], p. 408, $T_{X/S}$ is an invertible sheaf because it is a second syzygy of the $\mathcal{O}_X$ -module N .

Lemma 1.1 *If all the singular fibres of f are reduced then*

$$T_{X/S} \cong \omega_{X/S}^{-1} \quad (5)$$

Proof. Since all singular fibres of f are reduced, the locus where f is not smooth is finite, and therefore $\Omega_{X/S}^1$ and $\omega_{X/S}$ are isomorphic in codimension one. We thus have

$$c_1(\Omega_{X/S}^1) = c_1(\omega_{X/S})$$

and the conclusion follows because $T_{X/S}$ is invertible. $\square$

Remark 1.2 If f has some non-reduced singular fibre then $T_{X/S}$ and $\omega_{X/S}^{-1}$ are not isomorphic. Precisely we have:

$$T_{X/S} = \omega_{X/S}^{-1}(\sum (\nu_i - 1)E_i)$$

where $\{\nu_i E_i\}$ is the set of all components of the singular fibres of f , and where ν_i denote their corresponding multiplicities. This formula is due to Serrano ([25], Lemma 1.1).

Next proposition will not be applied in the sequel:

Proposition 1.3 *If f is any fibration we have:*

$$\chi(\omega_{X/S}^{-1}) = \chi(\mathcal{O}_X) + K_X^2 - 6(b-1)(g-1)$$

If f has reduced singular fibres then

$$\chi(T_{X/S}) = \chi(\mathcal{O}_X) + K_X^2 - 6(b-1)(g-1)$$

Proof. By Riemann–Roch:

$$\chi(\omega_{X/S}^{-1}) = \chi(\mathcal{O}_X) + \frac{1}{2}[\omega_{X/S} \cdot \omega_{X/S} + \omega_X \cdot \omega_{X/S}] =$$

(replacing $\omega_X \otimes f^*\omega_S^{-1}$ for $\omega_{X/S}$)

$$\begin{aligned} &= \chi(\mathcal{O}_X) + \frac{1}{2}[K_X^2 - 2(2b-2)(2g-2) + K_X^2 - (2b-2)(2g-2)] = \\ &= \chi(\mathcal{O}_X) + K_X^2 - 6(b-1)(g-1) \end{aligned}$$

The last formula is a consequence of Lemma 1.1. $\square$

The following is a classical result of Arakelov in the semistable case, and it is due to Serrano in the general case (recall that we are assuming $g \geq 2$):

Theorem 1.4 *If f is a non-isotrivial fibration then*

$$h^0(X, T_{X/S}) = 0 = h^0(X, T_X) \quad (6)$$

If moreover f is relatively minimal then we also have:

$$h^1(X, T_{X/S}) = 0 \quad (7)$$

Proof. f is non-isotrivial if and only if $f_*T_X = 0$ ([26], Lemma 3.2). Since

$$f_*T_{X/S} \subset f_*T_X$$

we also have $f_*T_{X/S} = 0$ if f is non-isotrivial. Thus (6) is a consequence of the Leray spectral sequence. For (7) see [1] or [27] in the semistable case, and [25], Corollary 3.6, in the general case. $\square$

Denoting by Ext_f^1 the first derived functor of f_*Hom , we are interested in the sheaf $Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X)$ because its cohomology describes the deformation theory of f (see §2).

Lemma 1.5 *For any fibration $f : X \rightarrow S$ there is an isomorphism*

$$Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X) \cong Hom(f_*(\Omega_{X/S}^1 \otimes \omega_{X/S}), \mathcal{O}_S) \quad (8)$$

so that in particular the first sheaf is locally free of rank $3g - 3$. Moreover there is an exact sequence of sheaves on S :

$$\begin{array}{ccccccc} 0 & \rightarrow & R^1 f_* T_{X/S} & \xrightarrow{c_{10}} & Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X) & \xrightarrow{c_{01}} & f_* Ext_X^1(\Omega_{X/S}^1, \mathcal{O}_X) \rightarrow 0 \\ & & & & & & \parallel \\ & & & & & & f_* N \end{array} \quad (9)$$

Proof. (8) is just relative duality for the fibration f ([18], Corollary (24)), because $\Omega_{X/S}^1$ is S -flat.

(9) is the sequence associated to the local-to-global spectral sequence for Ext_f . $\square$

Proposition 1.6 *If the fibration f is non-isotrivial then we have:*

$$\chi(Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X)) = 11\chi(\mathcal{O}_X) - 2K_X^2 + 2(b-1)(g-1) \quad (10)$$

Proof. Since the fibres of f are 1-dimensional we have

$$R^2 f_* T_{X/S} = 0$$

Moreover $f_* Ext_f^2(\Omega_{X/S}^1, \mathcal{O}_X) = 0$ because $Ext^2(\Omega_{X/S}^1, \mathcal{O}_X) = 0$ by the exact sequence (2). Therefore, using the local-to-global spectral sequence for Ext_f we deduce that

$$R^1 f_* N = Ext_f^2(\Omega_{X/S}^1, \mathcal{O}_X) = 0$$

where the last equality is true because the fibres of f are 1-dimensional. This gives:

$$\chi(f_* N) = \chi(N)$$

Moreover, since f is non-isotrivial, from (6) and the Leray spectral sequence we get

$$\chi(R^1 f_* T_{X/S}) = -\chi(T_{X/S})$$

We now use the exact sequence (9) and we deduce that

$$\begin{aligned} \chi(Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X)) &= \chi(f_* N) + \chi(R^1 f_* T_{X/S}) \\ &= \chi(N) - \chi(T_{X/S}) \\ &= \chi(f^* T_S) - \chi(T_X) \quad (\text{by (3)}) \end{aligned}$$

Using Riemann-Roch one computes that:

$$\begin{aligned}\chi(f^*(T_S)) &= \chi(\mathcal{O}_X) + 2(b-1)(g-1) \\ \chi(T_X) &= 2K_X^2 - 10\chi(\mathcal{O}_X)\end{aligned}$$

and by substitution one gets (10). $\square$

Assume that the fibration f is semistable. In this case N is a torsion sheaf supported on the set of double points of the fibres of f , and having stalk $\mathbf{k}$ at each of them. In particular we have

$$h^0(X, N) =: \delta \quad (11)$$

the total number of double points of the fibres of f . Moreover $\Omega_{X/S}^1$ and $\omega_{X/S}$ are related by the exact sequence ([16], p. 49):

$$0 \rightarrow \Omega_{X/S}^1 \rightarrow \omega_{X/S} \rightarrow N \rightarrow 0 \quad (12)$$

We will denote by $\mathcal{M}_g$ (respectively $\overline{\mathcal{M}}_g$) the moduli stack of nonsingular projective curves (respectively stable curves) of genus g . To the semistable fibration f there is associated a flat family of stable curves

$$f' : X' \rightarrow S$$

where X' is the surface obtained from X after contracting all the (-2) -curves in the fibres. Therefore we can associate to f' , hence to f , a modular morphism

$$\psi_f : S \rightarrow \overline{\mathcal{M}}_g$$

We know ([16], p. 149) that:

$$f_*(\Omega_{X/S}^1 \otimes \omega_{X/S}) = \psi_f^* \Omega_{\overline{\mathcal{M}}_g}^1$$

From (8) it follows that:

$$Ext_f^1(\Omega_{X/S}^1, \mathcal{O}_X) = \psi_f^* T_{\overline{\mathcal{M}}_g} \quad (13)$$

Proposition 1.7 *If the fibration f is semistable and non-isotrivial then we have:*

$$\chi(\psi_f^* T_{\overline{\mathcal{M}}_g}) = 11\chi(\mathcal{O}_X) - 2K_X^2 + 2(b-1)(g-1) \quad (14)$$

and

$$c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) = -5(b-1)(g-1) - [11\chi(\mathcal{O}_X) - 2K_X^2] \quad (15)$$

Proof. Identity (14) is a combination of (13) and of Proposition 1.6. The Riemann-Roch theorem for the vector bundle $\psi_f^* T_{\overline{\mathcal{M}}_g}$ gives

$$\chi(\psi_f^* T_{\overline{\mathcal{M}}_g}) = -c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) + (1-b)(3g-3)$$

Comparing with the expression (14) we obtain (15). $\square$

2 Deformations of fibrations

In this section we will analyze the deformation-theoretic information carried by a fibration $f : X \rightarrow S$. We have a homomorphism:

$$\kappa_f : T_S \rightarrow \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X)$$

arising as the coboundary in the long exact sequence of $f_* \text{Hom}(-, \mathcal{O}_X)$ of the exact sequence (2).

Lemma 2.1 *For general $p \in S$ the homomorphism κ_f induces a map:*

$$\kappa_f(p) : T_p S \rightarrow \text{Ext}_{X(p)}^1(\Omega_{X(p)}^1, \mathcal{O}_{X(p)})$$

which coincides with the Kodaira-Spencer map of the family f at p . f is non-isotrivial if and only if κ_f is injective.

Proof. $\kappa_f(p)$ is the composition

$$T_p S \rightarrow \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X)_p \otimes \mathbf{k}(p) \xrightarrow{\tau_p^1} \text{Ext}_{X(p)}^1(\Omega_{X(p)}^1, \mathcal{O}_{X(p)})$$

of the natural restriction of κ with the base change map τ_p^1 . If $X(p)$ is reduced then $\text{Ext}_{X(p)}^1(\Omega_{X(p)}^1, \mathcal{O}_{X(p)})$ is the vector space of first order deformations of $X(p)$ and $\kappa_f(p)$ is the Kodaira-Spencer map of f at p essentially by definition (see [24], Remark 2.4.4).

The exact sequence (2) induces an exact sequence on S :

$$0 \rightarrow f_* T_X \rightarrow T_S \xrightarrow{\kappa_f} \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X)$$

We have $f_* T_X = 0$ if and only if f is non-isotrivial, by Lemma 3.2 of [26], and this proves the last assertion. $\square$

Lemma 2.2 *Let $f : X \rightarrow S$ be a fibration, and let Def_f be the functor of Artin rings of deformations of f leaving the target fixed (see [24]). Then there is a natural isomorphism*

$$\mu : \text{Ext}_X^1(\Omega_{X/S}^1, \mathcal{O}_X) \rightarrow H^0(S, \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X))$$

and both spaces are naturally identified with the tangent space of Def_f . Moreover

$$H^1(S, \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X))$$

is an obstruction space for Def_f .

Proof. There is an exact sequence:

$$0 \rightarrow H^1(S, f_*T_{X/S}) \rightarrow \text{Ext}_X^1(\Omega_{X/S}^1, \mathcal{O}_X) \xrightarrow{\mu} H^0(S, \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X)) \rightarrow 0$$

([19], p. 105). Since $f_*T_{X/S} = 0$ (see the proof of Theorem 1.4) μ is an isomorphism. First order deformations of f are in 1-1 correspondence with isomorphism classes of extensions

$$\zeta : 0 \rightarrow \mathcal{O}_X \xrightarrow{j} \mathcal{O}_{\mathcal{X}} \rightarrow \mathcal{O}_X \rightarrow 0$$

such that $\mathcal{O}_{\mathcal{X}}$ is a sheaf of flat $\mathcal{O}_S[\epsilon]$ -algebras. In [24], p. 104, it is proved that if we only assume that ζ is an extension of $\mathcal{O}_S$ -algebras then it has already a structure of deformation by sending $\epsilon \mapsto j(1)$. There is a natural 1-1 correspondence between isomorphism classes of extensions ζ as above and $\text{Ext}_X^1(\Omega_{X/S}^1, \mathcal{O}_X)$ (loc. cit., Theorem 1.1.10).

By comparing the cohomology sequence of (9) with the exact sequence (d) of Lemma 3.4.7 in loc. cit., we see that $H^1(S, \text{Ext}_f^1(\Omega_{X/S}^1, \mathcal{O}_X))$ is an obstruction space for Def_f . $\square$

We now consider the case $S = \mathbb{P}^1$, i.e. a fibration

$$f : X \rightarrow \mathbb{P}^1$$

It will be called a *rational fibration*. We have:

$$\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \cong \bigoplus_{i=1}^{3g-3} \mathcal{O}_{\mathbb{P}^1}(a_i)$$

for some integers a_i . We call f *free* (resp. *very free*) if it is non-isotrivial and $a_i \geq 0$ for all i (resp. if it is non-isotrivial and $a_i \geq 1$ for all i).

Next proposition gives necessary numerical conditions for a rational fibration to be free.

Proposition 2.3 *Suppose that $f : X \rightarrow \mathbb{P}^1$ is a rational fibration. If f is free then we have:*

$$11\chi(\mathcal{O}_X) - 2K_X^2 \geq 5(g-1) + 2 \quad (16)$$

If f is very free then:

$$11\chi(\mathcal{O}_X) - 2K_X^2 \geq 8(g-1) + 1 \quad (17)$$

Proof. Since f is non-isotrivial the homomorphism

$$\kappa_f : T_{\mathbb{P}^1} \rightarrow \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)$$

induced by the exact sequence (2) is injective, by Lemma 2.1. It follows that

$$\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \cong \bigoplus_{i=1}^{3g-3} \mathcal{O}_{\mathbb{P}^1}(a_i)$$

with $a_i \geq 2$ for some i . We thus have

$$h^0(\mathbb{P}^1, \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \geq 3g-1$$

if f is free and

$$h^0(\mathbb{P}^1, \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \geq 6g-5$$

if f is very free. Therefore

$$\begin{aligned} 11\chi(\mathcal{O}_X) - 2K_X^2 - 2(g-1) &= \chi(\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \\ &= h^0(\mathbb{P}^1, \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \\ &\geq \begin{cases} 3g-1 & \text{if } f \text{ free} \\ 6g-5 & \text{if } f \text{ very free} \end{cases} \end{aligned}$$

where the first equality is Proposition 1.6. $\square$

Definition 2.4 *Let $f : X \rightarrow \mathbb{P}^1$ be a non-isotrivial rational fibration. We say that the fibres of f have general moduli if there is a family of deformations of f :*

$$\begin{array}{ccc} X & \xrightarrow{\quad \quad} & \mathcal{X} \\ \downarrow f & & \downarrow F \\ \mathbb{P}^1 \times \{v\} & \xrightarrow{\quad \quad} & \mathbb{P}^1 \times V \end{array} \quad \begin{array}{c} \nearrow \varphi \\ \nearrow \end{array} \quad \begin{array}{c} V \\ \end{array}$$

parametrized by a nonsingular connected pointed algebraic scheme (V, v) such that F is a family of curves of genus g with surjective Kodaira-Spencer map at every point of a non-empty open subset of $\mathbb{P}^1 \times \{v\}$.

From the definition it follows that if the fibres of f have general moduli then, up to taking a general deformation of f , we may assume that a general fibre C of f is a general curve of genus g . The following are some well known properties of a general curve which we will use:

Proposition 2.5 *Let C be a general nonsingular connected projective curve of genus $g \geq 2$. Then:*

(i) C has Clifford index

$$\text{Cliff}(C) = \left\lfloor \frac{g-1}{2} \right\rfloor$$

(ii) For any invertible sheaf L on C , of degree $d \geq 0$ and with $h^0(L) = r+1 \geq 1$ we have

$$\rho(g, r, d) \geq 0$$

where $\rho(g, r, d) := g - (r+1)(g-d+r)$ is the Brill-Noether number. In particular:

$$d \geq \frac{1}{2}g + 1 \quad \text{if } r \geq 1$$

$$d \geq \frac{2}{3}g + 2 \quad \text{if } r \geq 2$$

(iii) $H^1(L^2) = 0$ for any invertible sheaf L on C such that $h^0(L) \geq 2$.

(iv) C does not possess irrational involutions, i.e. non-constant morphisms of degree ≥ 2 onto a curve of positive genus.

Proof. (i) follows from the main result of [10] and from Brill-Noether theory. (ii) is proved in [15]. For (iii) see [5], p. 22, and references therein. (iv) is classical and follows from Riemann's existence theorem and a count of constants. $\square$

Definition 2.4 is related with the notion of free rational fibration.

Theorem 2.6 *Assume that $f : X \rightarrow \mathbb{P}^1$ is a non-isotrivial rational fibration whose fibres have general moduli. Then f is free.*

Proof. By assumption there is a family of deformations of f as in Definition 2.4. We have an exact sequence of sheaves on $\mathcal{X}$:

$$0 \rightarrow F^* \Omega_{\mathbb{P}^1 \times V}^1 \rightarrow \Omega_{\mathcal{X}}^1 \rightarrow \Omega_{\mathcal{X}/\mathbb{P}^1 \times V}^1 \rightarrow 0$$

Taking $F_* \text{Hom}(-, \mathcal{O}_{\mathcal{X}})$ we obtain a coboundary homomorphism of sheaves on $\mathbb{P}^1 \times V$:

$$\kappa : T_{\mathbb{P}^1 \times V} \rightarrow \text{Ext}_f^1(\Omega_{\mathcal{X}/\mathbb{P}^1 \times V}^1, \mathcal{O}_{\mathcal{X}})$$

For a general $p \in \mathbb{P}^1$ the composition:

$$T_{(p,v)}(\mathbb{P}^1 \times V) \xrightarrow{\kappa(p,v)} \text{Ext}_f^1(\Omega_{\mathcal{X}/\mathbb{P}^1 \times V}^1, \mathcal{O}_{\mathcal{X}})_{(p,v)} \otimes \mathbf{k} \xrightarrow{\tau_{(p,v)}^1} \text{Ext}^1(\Omega_{X(p)}^1, \mathcal{O}_{X(b)}) \quad (18)$$

is the Kodaira-Spencer map of F at (p, v) , and by hypothesis this is surjective.

Restricting to $\mathbb{P}^1 \times \{v\}$ we obtain:

$$(T_{\mathbb{P}^1 \times V}) \otimes \mathcal{O}_{\mathbb{P}^1 \times \{v\}} \cong \mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}^{\dim(V)}$$

and a base change homomorphism:

$$\text{Ext}_f^1(\Omega_{\mathcal{X}/\mathbb{P}^1 \times V}^1, \mathcal{O}_{\mathcal{X}}) \otimes \mathcal{O}_{\mathbb{P}^1 \times \{v\}} \rightarrow \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)$$

which is an isomorphism on an open set because $\dim[\text{Ext}^1(\Omega_{X(p)}^1, \mathcal{O}_{X(b)})] = 3g - 3$ for all p in an open set of $\mathbb{P}^1$. Therefore we obtain a homomorphism of locally free sheaves on $\mathbb{P}^1$:

$$\mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}_{\mathbb{P}^1}^{\dim(V)} \rightarrow \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \quad (19)$$

Combining with (18) and using the fact that $\tau_{(p,v)}^1$ is an isomorphism on an open set, we obtain that (19) is surjective on an open set. This implies that $\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)$ is globally generated, or, what is the same, that f is free. $\square$

Typical examples of rational fibrations are obtained as follows. Let Y be a projective nonsingular surface, and let $C \subset Y$ be a projective nonsingular connected curve of genus g such that

$$\dim(|C|) \geq 1$$

Consider a linear pencil Λ contained in $|C|$ whose general member is nonsingular and let $\sigma : X \rightarrow Y$ be the blow-up at its base points. We obtain a rational fibration

$$f : X \rightarrow \mathbb{P}^1$$

obtained by composing σ with the rational map $Y \dashrightarrow \mathbb{P}^1$ defined by Λ . We will call f the fibration defined by the pencil Λ .

If we assume that all singular elements of a general linear pencil in $|C|$ are nodal, then a general linear pencil contained in $|C|$ defines a fibration which becomes semistable after contracting the (-1) -curves contained in the fibres.

Corollary 2.7 *Assume that a general projective nonsingular connected curve C of genus $g \geq 3$ can be embedded in a projective nonsingular non-ruled surface Y as a member of a non-trivial linear system. Then a general pencil $\Lambda \subset |C|$ containing C as a member defines a free fibration.*

Proof. Let $f : X \rightarrow \mathbb{P}^1$ be the fibration defined by the pencil Λ .

Assume that f is isotrivial. Then, by the structure theorem for isotrivial fibrations [26], X is birational to a quotient $(C \times B)/G$, where B is a nonsingular curve and G is a finite group acting on both C and B . But since C is a general curve of genus g , it has no non-trivial automorphisms: this implies that G acts trivially on C and it follows that X is birational to $C \times \mathbb{P}^1$. We have a contradiction because Y is not ruled, and therefore f is non-isotrivial.

By hypothesis there is a pointed nonsingular algebraic scheme (V, v) and a commutative diagram as follows:

$$\begin{array}{ccccc} C & \hookrightarrow & \mathcal{C} & \hookrightarrow & \mathcal{Y} \\ \downarrow & & \downarrow & \nearrow \beta & \\ \mathrm{Spec}(\mathbf{k}) & \xrightarrow{v} & V & & \end{array}$$

such that the left square is a smooth family of deformations of C having surjective Kodaira-Spencer map at every point; β is a smooth family of projective surfaces and the upper right inclusion restricts over v to the inclusion $C \subset Y$; moreover $|\mathcal{C}(w)|$ is a non-trivial linear system on the surface $\mathcal{Y}(w)$ for all closed points $w \in V$.

Let $\mathcal{L} = \mathcal{O}_{\mathcal{Y}}(\mathcal{C})$. After possibly performing an etale base change, we can find a trivial free subsheaf of rank two $\mathcal{O}_V^{\oplus 2} \subset \beta_* \mathcal{L}$ which defines a rational V -map

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{\psi} & \mathbb{P}^1 \times V \\ \downarrow \beta & \nearrow & \\ V & & \end{array} \quad (20)$$

whose restriction over v is the rational map defined by the pencil Λ . Let $Z \subset \mathcal{Y}$ be the scheme of indeterminacy of ψ and let $\theta : \mathcal{X} \rightarrow \mathcal{Y}$ be the blow-up with center Z . Composing with θ we obtain from (20) a family of deformations of f :

$$\begin{array}{ccccc}
 X & \hookrightarrow & \mathcal{X} & & \\
 \downarrow f & & \downarrow \psi \cdot \theta & \searrow \beta \cdot \theta & \\
 \mathbb{P}^1 \times \{v\} & \hookrightarrow & \mathbb{P}^1 \times V & \nearrow & V
 \end{array}$$

By construction this diagram implies that the fibres of f have general moduli, and the corollary follows from Theorem 2.6. $\square$

We have the following remark concerning non-rational ruled surfaces.

Proposition 2.8 *Assume that a general projective nonsingular connected curve C of genus $g \geq 3$ can be embedded in a projective nonsingular non-rational birationally ruled surface Y . Then Y is birationally equivalent to $C \times \mathbb{P}^1$. Moreover any linear pencil in $|C|$ containing C as a member defines an isotrivial fibration.*

Proof. By hypothesis there is a birational isomorphism $\xi : Y \dashrightarrow \Gamma \times \mathbb{P}^1$ for some projective nonsingular curve Γ of positive genus. The composition

$$h : C \hookrightarrow Y \xrightarrow{\xi} \Gamma \times \mathbb{P}^1 \longrightarrow \Gamma$$

where the last morphism is the projection, is non-constant. Since C is a general curve, it does not possess irrational involutions (Prop. 2.5(iv)), thus h must be an isomorphism. The last assertion of the proposition follows from the fact that any general member D of the pencil is again a general curve of genus g and therefore $D \cong \Gamma \cong C$. $\square$

3 Rational fibrations defined by linear pencils

In this section we will consider a pencil in a linear system $|C|$ on a surface Y and we will rephrase the numerical conditions given in §2 in terms of invariants of Y and of the linear system $|C|$.

Lemma 3.1 *Let Y be a projective nonsingular surface, $C \subset Y$ a projective nonsingular connected curve of genus g such that*

$$r := \dim(|C|) \geq 1$$

Let $f : X \rightarrow \mathbb{P}^1$ be the fibration defined by a general pencil contained in $|C|$, and assume that it is non-isotrivial. Then we have an inclusion

$$\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(r-1)} \subset \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)$$

Proof. Let

$$\begin{array}{ccc} \mathcal{X} & \hookrightarrow & \mathbb{P}^r \times Y \\ \downarrow F & & \\ \mathbb{P}^r & & \end{array}$$

be the family of curves in Y parametrized by the linear system $|C| \cong \mathbb{P}^r$. We have an exact sequence:

$$0 \rightarrow F^*\Omega_{\mathbb{P}^r}^1 \rightarrow \Omega_{\mathcal{X}}^1 \rightarrow \Omega_{\mathcal{X}/\mathbb{P}^r}^1 \rightarrow 0$$

Taking $F_*\text{Hom}(-, \mathcal{O}_{\mathcal{X}})$ we obtain an exact sequence:

$$F_*T_{\mathcal{X}} \rightarrow T_{\mathbb{P}^r} \rightarrow \text{Ext}_F^1(\Omega_{\mathcal{X}/\mathbb{P}^r}^1, \mathcal{O}_{\mathcal{X}})$$

Denote by

$$\zeta : \mathbb{P}^1 \rightarrow \mathbb{P}^r$$

the line corresponding to the pencil defining f . Restriction to the line ζ induces the following commutative diagram:

$$\begin{array}{ccccc} \zeta^*F_*T_{\mathcal{X}} & \rightarrow & \zeta^*T_{\mathbb{P}^r} & \rightarrow & \zeta^*\text{Ext}_F^1(\Omega_{\mathcal{X}/\mathbb{P}^r}^1, \mathcal{O}_{\mathcal{X}}) \\ \downarrow & & & & \downarrow \\ f_*T_X & \rightarrow & T_{\mathbb{P}^1} & \xrightarrow{\kappa_f} & \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \end{array}$$

Since f is non-isotrivial, $f_*T_X = 0$ by Lemma 3.2 of [26], and both vertical maps are isomorphisms at a general point $p \in \mathbb{P}^1$. This implies that $\zeta^*T_{\mathbb{P}^r}$ is a subsheaf of $\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)$. It follows that, up to order, we have

$$a_1 \geq 2, \quad a_i \geq 1, \quad i = 2, \dots, r$$

because $\zeta^*T_{\mathbb{P}^r} \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(r-1)}$ ([2], Example 2.7). $\square$

Theorem 3.2 *Let Y be a projective nonsingular surface, $C \subset Y$ a projective nonsingular connected curve of genus g such that*

$$\dim(|C|) \geq 1$$

Let $f : X \rightarrow \mathbb{P}^1$ be the fibration defined by a general pencil contained in $|C|$. Then:

(i) *If f is free we have:*

$$5(g-1) - [11\chi(\mathcal{O}_Y) - 2K_Y^2 + 2C^2] + h^0(C) \leq 0 \quad (21)$$

If moreover $h^0(K_Y - C) = 0$ then

$$4(g-1) - C^2 - [10\chi(\mathcal{O}_Y) - 2K_Y^2] \leq 0 \quad (22)$$

(ii) *If f is very free then*

$$8(g-1) - [11\chi(\mathcal{O}_Y) - 2K_Y^2 + 2C^2] + h^0(C) \leq 0 \quad (23)$$

If moreover $h^0(K_Y - C) = 0$ then

$$7(g-1) - C^2 - [10\chi(\mathcal{O}_Y) - 2K_Y^2] \leq 0 \quad (24)$$

Proof. Letting $d := C^2$, we have $K_X^2 = K_Y^2 - d$.

Assume that f is free. From Proposition 1.6 we derive in this case:

$$\begin{aligned} h^0(\mathbb{P}^1, \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) &= \chi(\text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \\ &= 11\chi(\mathcal{O}_Y) - 2K_Y^2 + 2d - 2(g-1) \end{aligned} \quad (25)$$

From Lemma 3.1 we have:

$$3g - 3 + h^0(C) \leq h^0(\mathbb{P}^1, \text{Ext}_f^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X)) \quad (26)$$

Putting (25) and (26) together we obtain (21).

Applying the Riemann–Roch theorem we have:

$$\chi(C) = \chi(\mathcal{O}_Y) - (g-1-d)$$

and therefore:

$$h^0(C) \geq \chi(\mathcal{O}_Y) - (g-1-d)$$

provided $h^0(K_Y - C) = 0$. Substituting in (21) we get (22).

(ii) is proved similarly. $\square$

Proposition 3.3 *In the situation of Theorem 3.2 assume moreover that the fibration f is semistable. Letting $\psi_f : \mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_g$ be the modular morphism defined by f , have:*

$$c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) = 5(g-1) - [11\chi(\mathcal{O}_Y) - 2K_Y^2 + 2C^2] \quad (27)$$

and condition (21) is equivalent to:

$$c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) + h^0(C) \leq 0 \quad (28)$$

Proof. Since $K_X^2 = K_Y^2 - C^2$ and $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_Y)$, comparing with (15) we obtain (27). The equivalence between (21) and (28) is obvious. $\square$

Remarks 3.4 (i) Recall that if $h^0(T_Y) = 0$ then

$$10\chi(\mathcal{O}_Y) - 2K_Y^2 = -\chi(T_Y) = h^1(T_Y) - h^2(T_Y) =: \mu(Y)$$

the expected number of moduli of the surface Y (see [24]). Inequalities (21),..., (24) of Theorem 3.2 say that $\mu(Y)$ must be sufficiently high with respect to g in order for Y to contain pencils of curves of genus g defining free or very free semistable fibrations.

(ii) In [9] it is shown that a general curve C of genus 15 can be embedded as a non-degenerate nonsingular curve in $\mathbb{P}^6$, lying in a nonsingular canonical surface $Y \subset \mathbb{P}^6$ complete intersection of 4 quadrics, and that $\dim(|C|) = 2$ on Y . Therefore, by Theorem 2.6, a general pencil in $|C|$ defines a free rational fibration. The relevant numbers are in this case:

$$d = C^2 = 9, \quad K_Y^2 = 16, \quad \chi(\mathcal{O}_Y) = 8$$

We find:

$$10\chi(\mathcal{O}_Y) - 2K_Y^2 = 48 > 47 = 4(g-1) - d$$

as it must be, because of the existence of a free rational fibration of genus 15. But

$$48 < 89 = 7(g-1) - d$$

which shows that such a fibration is not very free.

So far we have considered, on a surface Y , linear pencils whose general member is a nonsingular curve. It is not difficult to obtain an estimate analogous to (22) which applies to linear pencils whose general member is an irreducible *nodal* curve.

Proposition 3.5 *Let Y be a projective nonsingular surface, and let $|C|$ be a linear system whose general member is irreducible of arithmetic genus g . Let*

$$\Lambda \subset |C|$$

be a linear pencil consisting of curves of $|C|$ which are singular at given distinct points $p_1, \dots, p_\delta \in Y$. Assume the following:

- (i) *The general member of Λ is irreducible and has $p_1, \dots, p_\delta$ as nodes and no other singularities, so that it has geometric genus $\gamma := g - \delta \geq 2$.*
- (ii) *$h^0(K_Y - C) = 0$.*

Let $f : X \rightarrow \mathbb{P}^1$ be the rational fibration associated to the proper transform of Λ on the surface Z obtained from Y after blowing-up $p_1, \dots, p_\delta$.

If f is a free fibration then

$$4(g - 1) - C^2 - [10\chi(\mathcal{O}_Y) - 2K_Y^2] \leq 2\delta \quad (29)$$

or, equivalently:

$$4(\gamma - 1) - C^2 - [10\chi(\mathcal{O}_Y) - 2K_Y^2] \leq -2\delta \quad (30)$$

Proof. It is not restrictive to assume that C is a general element of Λ ; let $\tilde{C} \subset Z$ be the proper transform of C . Then $\tilde{C}$ has genus $\gamma := g - \delta$, and $\tilde{C}^2 \leq C^2 - 4\delta$. Moreover assumption (ii) implies that $h^0(K_Z - \tilde{C}) = 0$. By applying (22) to the proper transform of Λ on Z we obtain:

$$\begin{aligned} 0 &\geq 4(\gamma - 1) - \tilde{C}^2 - [10\chi(\mathcal{O}_Z) - 2K_Z^2] \\ &\geq 4(g - 1) - 4\delta - (C^2 - 4\delta) - [10\chi(\mathcal{O}_Y) - 2(K_Y^2 - \delta)] \\ &= 4(g - 1) - C^2 - [10\chi(\mathcal{O}_Y) - 2K_Y^2] - 2\delta \end{aligned}$$

The equivalence of (29) and (30) is obvious. □

4 Curves on rational surfaces

In this section we consider linear pencils of nodal curves on a minimal rational surface. We start from the case of $\mathbb{P}^2$.

Theorem 4.1 *Let $C \subset \mathbb{P}^2$ be an irreducible curve of degree n having δ nodes $p_1, \dots, p_\delta$ and no other singularities. Let*

$$\gamma = \binom{n-1}{2} - \delta$$

be the geometric genus of C . Assume that:

- (i) There exists a linear pencil of curves of degree n singular at $p_1, \dots, p_\delta$.*
- (ii) $n \geq \frac{2}{3}\gamma + 2$.*

Then $n \leq 8$ and $\gamma \leq 9$.

The same conclusion holds if we replace (ii) by the assumption that the normalization $\tilde{C}$ of C is a general curve of genus γ .

Proof. (i) implies that on $\tilde{C}$ there is an *effective* divisor of degree $n^2 - 4\delta$. On the other hand (ii) implies that

$$3n \geq n^2 - 3n + 2 - 2\delta + 6$$

equivalently:

$$2\delta \geq n^2 - 6n + 8$$

Therefore:

$$0 \leq n^2 - 4\delta \leq -n^2 + 12n - 16$$

which implies $n \leq 10$. An elementary well known case by case analysis shows that there are no nodal irreducible curves of degree 9 or 10 satisfying both (i) and (ii), and that in degree $n = 8$ the highest genus of a nodal curve satisfying (i) and (ii) is $\gamma = 9$, corresponding to $\delta = 12$ (see [6] for details).

If $\tilde{C}$ is a general curve of genus γ then inequality (ii) follows from Proposition 2.5(ii) applied to $L = \mathcal{O}_{\tilde{C}}(1)$. Therefore the last assertion follows from the first part. $\square$

Let $Y = F_n := \mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(n) \oplus \mathcal{O}_{\mathbb{P}^1})$, $n \geq 0$, $n \neq 1$, be a minimal rational ruled surface. Denote by $F \subset F_n$ a fibre of the projection $F_n \rightarrow \mathbb{P}^1$, and

by $E \subset F_n$ a curve of self-intersection $E^2 = n$ such that $EF = 1$. Then for every irreducible nonsingular curve $C \subset F_n$ we have

$$C \sim \alpha E + \beta F$$

for some $\alpha, \beta \geq 0$.

Theorem 4.2 *Keeping the above notations, let $C \sim \alpha E + \beta F$, $\alpha, \beta > 0$, be an irreducible curve on F_n having δ nodes $p_1, \dots, p_\delta$ and no other singularities, and geometric genus $\gamma \geq 2$. Assume that:*

- (i) *there exists a linear pencil $\Lambda \subset |C|$ such that $C \in \Lambda$ and all members of Λ are singular at $p_1, \dots, p_\delta$.*
- (ii) *The normalization $\tilde{C}$ of C is a general curve of genus γ .*

Then $\gamma \leq 14$.

Proof. The assumptions imply that a general member of Λ is a nodal curve. A general member of $|C|$ is irreducible and nonsingular of genus

$$g = \frac{1}{2}(\alpha - 1)(\alpha n + 2\beta - 2)$$

so that

$$\gamma = g - \delta$$

Moreover we have:

$$C^2 = 2\alpha\beta + n\alpha^2$$

Let $\varphi : \tilde{C} \rightarrow C \subset F_n$ be the composition of the normalization of C with the inclusion. The existence of Λ implies that

$$0 \leq \deg(\varphi^* \mathcal{O}(C)) - 4\delta = C^2 - 4\delta = 2\alpha\beta + n\alpha^2 - 4\delta \quad (31)$$

On the other hand, using Corollary 2.7 we see that the hypotheses of Proposition 3.5 are satisfied. In this case it can be easily computed that inequality (29) takes the following form:

$$2\delta + 2 \geq (\alpha - 2)(n\alpha + 2\beta - 4) \quad (32)$$

Putting (31) and (32) together we obtain:

$$0 \leq 2\alpha\beta + n\alpha^2 - 2(\alpha - 2)(n\alpha + 2\beta - 4) + 4$$

which is equivalent to:

$$0 \leq -n\alpha^2 + (4n - 2\beta + 8)\alpha + 8\beta - 12$$

This inequality implies that:

- if $n = 0$ then either $\alpha \leq 8$ or $\beta \leq 8$;
- if $n \geq 2$ then $\alpha \leq 8$.

In case $n = 0$ assume for simplicity that $\alpha \leq 8$. Since the composition

$$\tilde{C} \xrightarrow{\varphi} F_n \xrightarrow{\pi} \mathbb{P}^1$$

defines a g_α^1 on $\tilde{C}$, which is a general curve of genus γ , it follows (Proposition 2.5(ii)) that

$$\gamma \leq 2\alpha - 2 \leq 14$$

□

Remark 4.3 I don't know whether the bound of Theorem 4.2 is sharp.

5 Curves on surfaces of small Kodaira dimension

In this section we will prove some results showing that the existence of a curve with general moduli moving in a non-trivial linear system on a minimal surface Y such that $K_Y^2 = 0$ or having small geometric genus, imposes strong restrictions on the genus of the curve. Our analysis will cover all cases of surfaces with Kodaira dimension 0 or 1.

The following theorem is a simple consequence of Proposition 3.5. It gives bounds which, although not sharp, will be sufficient for our purposes.

Theorem 5.1 *Let Y be a minimal surface of geometric genus p_g and with $k\text{-dim}(Y) \geq 0$, and let $C \subset Y$ be an irreducible curve of geometric genus $\gamma \geq 3$ having nodes $p_1, \dots, p_\delta$ and no other singularities. Assume that:*

- (i) *C is contained in a linear pencil Λ whose general member is an irreducible curve with nodes at $p_1, \dots, p_\delta$ and no other singularities.*

(ii) The normalization $\tilde{C}$ of C is a general nonsingular curve of genus γ .

(iii) $h^0(K_Y - C) = 0$.

Then

$$\gamma \leq 1 + 5\chi(\mathcal{O}_Y) - K_Y^2$$

In particular

$$\gamma \leq \begin{cases} 6 & \text{if } p_g = 0 \\ 11 & \text{if } p_g = 1 \\ 16 & \text{if } p_g = 2 \end{cases}$$

Proof. C has arithmetic genus $g = \gamma + \delta$. By Corollary 2.7 applied to $\tilde{C} \subset Z$, where Z is the blow-up of Y at $p_1, \dots, p_\delta$, we may assume that the proper transform of Λ defines a free fibration. Therefore Proposition 3.5 applies. Since Y is not ruled, we have $K_Y^2 \geq 0$; moreover $\omega_Y^{\otimes n}$ is globally generated for some $n \gg 0$ and therefore $C^2 \leq 2g - 2$. From (29) we get:

$$2(g - 1) \leq 4(g - 1) - C^2 \leq 10\chi(\mathcal{O}_Y) - 2K_Y^2 + 2\delta$$

and we obtain the stated inequalities. $\square$

Remarks 5.2 The bounds of Theorem 5.1 are not sharp if the surface Y is irregular. For example, in the case of an abelian surface Y , a more accurate computation would give the bound $\gamma \leq 1$.

The bound $g \leq 11$ for the genus of a *nonsingular* curve with general moduli on a K3 surface can be deduced from a count of parameters, as shown in [20]. In [21] and [22] it is proved that a general nonsingular curve of genus g can be embedded in a K3 surface if and only if $g \leq 11$ and $g \neq 10$. In particular the bound of Theorem 5.1 is sharp for K3 surfaces. In [14] it is shown that even a general curve of genus 10 can be birationally embedded in a K3 surface provided one allows it to have one node (hence arithmetic genus 11).

Note that, for a linear pencil generically consisting of nonsingular curves of genus g defining a semistable fibration on a K3 surface we have, by (27):

$$c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) = g - 23 \tag{33}$$

This means that $\psi_f^* K_{\overline{\mathcal{M}}_g}$ is non-negative when $g \geq 23$.

Similarly we obtain:

$$c_1(\psi_f^* K_{\overline{\mathcal{M}}_g}) = g - 12$$

for a linear pencil generically consisting of nonsingular curves of genus g defining a semistable fibration on an Enriques surface.

We now consider the case of minimal elliptic surfaces. Theorem 5.1 implies the bound $\gamma \leq 16$ for a general nodal curve moving in a pencil on such a surface provided $p_g \leq 2$. In the next theorem we confirm the same bound for higher values of p_g as well.

Theorem 5.3 *Let Y be a non-ruled and non-rational minimal elliptic surface of geometric genus p_g , $\pi : Y \rightarrow B$ the elliptic fibration onto a nonsingular connected curve B . Let $C \subset Y$ be an irreducible curve of geometric genus $\gamma \geq 3$ having nodes $p_1, \dots, p_\delta$ and no other singularities. Assume that:*

- (i) *C is contained in a linear pencil Λ whose general member is an irreducible curve with nodes at $p_1, \dots, p_\delta$ and no other singularities.*
- (ii) *The normalization $\tilde{C}$ of C is a general nonsingular curve of genus γ .*

Then $\gamma \leq 16$.

Proof. Let $|K| = |M| + D$, where $|M|$ is the mobile part and D is the fixed divisor. Then $M = \sum_{i=1}^m F_{b_i}$, where $b_1, \dots, b_m \in B$ and $F_{b_i} = \pi^{-1}(b_i)$. It follows that $h^0(K_Y - C) = 0$ because all fibres of π have arithmetic genus 1. From Theorem 5.1 we deduce that $\gamma \leq 16$ if $p_g \leq 2$. Therefore we may assume that $p_g \geq 3$.

Let F be a general fibre of π . We have $C \cdot F =: k \geq 1$. If $k = 1$ then $\pi|_C : C \rightarrow B$ is an isomorphism, and it follows that every linear pencil Λ containing C is isotrivial. Since C is general, it follows that Y is ruled (as in the proof of Corollary 2.7), which is a contradiction.

Therefore $k \geq 2$. Let $b : Z \rightarrow Y$ be the blow-up of Y at the points $p_1, \dots, p_\delta$, and let $E_i = b^{-1}(p_i)$, $i = 1, \dots, \delta$ be the exceptional curves. We have the following diagram:

$$\begin{array}{ccc} \tilde{C} & \xrightarrow{\tilde{j}} & Z \\ \downarrow \nu & & \downarrow b \\ C & \xrightarrow{j} & Y \\ & & \downarrow \pi \\ & & B \end{array}$$

where ν is the normalization morphism and $\tilde{j}$ and j are the inclusions. If B has genus $g(B) \geq 1$ then

$$\pi \circ b \circ \tilde{j} : \tilde{C} \rightarrow B$$

is an irrational involution on $\tilde{C}$, and this contradicts the fact that $\tilde{C}$ has general moduli (Proposition 2.5(iv)). Therefore $B \cong \mathbb{P}^1$, and $\pi \circ b \circ \tilde{j}$ defines a g_k^1 on $\tilde{C}$; in other words $h^0(\tilde{C}, \mathcal{O}_{\tilde{C}}(b^*F)) \geq 2$. Since $p_g \geq 3$, we have $m \geq 2$, and it follows that

$$\mathcal{O}_{\tilde{C}}(m \cdot b^*F) \subset \mathcal{O}_{\tilde{C}}(b^*K_Y) \subset \mathcal{O}_{\tilde{C}}(b^*K_Y + \sum E_i + \tilde{C}) = \omega_{\tilde{C}}$$

This means that $h^1(\tilde{C}, \mathcal{O}_{\tilde{C}}(m \cdot b^*F)) \neq 0$, while $h^0(\tilde{C}, \mathcal{O}_{\tilde{C}}(b^*F)) \geq 2$, contradicting the generality of $\tilde{C}$ (Proposition 2.5(iii)). $\square$

6 Curves on surfaces of general type

In this section we consider the case of curves on a minimal surface of general type Y . In case Y has geometric genus $p_g \leq 1$, Theorem 5.1 gives the bound $\gamma \leq 11$ for the geometric genus of a nodal irreducible curve with general normalization moving in a linear pencil on Y . In order to obtain significant bounds for all values of p_g we will need to assume that the nodal curve either moves in a larger linear system generically consisting of nodal curves of the same geometric genus, or is not contained in a canonical divisor. We will distinguish several cases according to the invariants of the surface.

Theorem 6.1 *Let Y be a minimal surface of general type. Assume that $C \subset Y$ is an irreducible curve of geometric genus $\gamma \geq 3$ having nodes $p_1, \dots, p_\delta$ and no other singularities. Denote by $|C(-2p_1 - \dots - 2p_\delta)|$ the linear subsystem of $|C|$ consisting of the effective divisors linearly equivalent to C and singular at $p_1, \dots, p_\delta$. Assume that the normalization $\tilde{C}$ of C is a general curve of genus γ . Then*

- (i) *If $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$ and $K_Y^2 \geq 3\chi(\mathcal{O}_Y) - 10$ then $\gamma \leq 19$.*
- (ii) *If $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$ and $K_Y^2 \geq 4\chi(\mathcal{O}_Y) - 16$ then $\gamma \leq 15$.*
- (iii) *If $\dim(|C(-2p_1 - \dots - 2p_\delta)|) = 1$, $|K_Y - C| = \emptyset$ and $K_Y^2 \geq 4\chi(\mathcal{O}_Y) - 16$ then $\gamma \leq 19$.*

Proof. By Theorem 5.1 we may assume that Y has geometric genus $p_g \geq 2$. C has arithmetic genus $g = \gamma + \delta$. Let $b : Z \rightarrow Y$ be the blow-up at $p_1, \dots, p_\delta$, and let $E_1, \dots, E_\delta \subset Z$ be the exceptional curves. Then we have the commutative diagram:

$$\begin{array}{ccc} \tilde{C} & \hookrightarrow & Z \\ \downarrow \nu & & \downarrow b \\ C & \hookrightarrow & Y \end{array}$$

where $\tilde{C} \sim b^*C - 2 \sum E_i$. Consider the line bundle $L := \mathcal{O}_{\tilde{C}}(b^*K_Y)$ on $\tilde{C}$. Then

$$h^0(\omega_{\tilde{C}}L^{-1}) = h^0(\mathcal{O}_{\tilde{C}}(\tilde{C} + \sum E_i)) \geq h^0(\mathcal{O}_{\tilde{C}}(\tilde{C}) \geq \dim(|C(-2p_1 - \dots - 2p_\delta)|))$$

Assume first that $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$. Then, by the previous inequality:

$$h^0(\omega_{\tilde{C}}L^{-1}) \geq 2 \quad (34)$$

If $h^0(Z, b^*K_Y - \tilde{C}) \neq 0$ then

$$2\tilde{C} < b^*K_Y + \tilde{C} < K_Z + \tilde{C}$$

so that $h^1(\tilde{C}, \mathcal{O}_{\tilde{C}}(2\tilde{C})) \neq 0$, and this is impossible because $\tilde{C}$ has general moduli (Proposition 2.5(iii)). Therefore $h^0(Z, b^*K_Y - \tilde{C}) = 0$. It follows that

$$h^0(L) \geq h^0(Z, b^*K_Y) = p_g \geq 2 \quad (35)$$

Inequalities (34) and (35) imply that L contributes to the Clifford index of $\tilde{C}$. Since $\tilde{C}$ has general moduli, recalling that

$$\text{Cliff}(L) = \deg(L) - 2h^0(L) + 2$$

we have the following relations satisfied by the Clifford indices:

$$\left\lfloor \frac{\gamma - 1}{2} \right\rfloor = \text{Cliff}(\tilde{C}) \leq \text{Cliff}(L) \leq \deg(L) - 2p_g + 2 \quad (36)$$

For the equality see Proposition 2.5(i). The first inequality follows from the very definition of $\text{Cliff}(\tilde{C})$, and the last is by (35).

Observe also that from the equality $h^0(Z, b^*K_Y - \tilde{C}) = 0$ it follows that $h^0(Y, K_Y - C) = 0$ as well. Moreover, by Theorem 2.6, a general linear pencil in $|\tilde{C}|$ defines a free fibration. Therefore Proposition 3.5 applies.

Let's prove (i). After substitution of the inequality $K_Y^2 \geq 3\chi(\mathcal{O}_Y) - 10$ in (30) we obtain:

$$4\chi(\mathcal{O}_Y) + 20 - 2\delta \geq 4(\gamma - 1) - C^2$$

which, using the inequality $\chi(\mathcal{O}_Y) \leq 1 + p_g$, can be replaced by the weaker

$$24 \geq 4(\gamma - 1) - C^2 - 4p_g + 2\delta$$

Since

$$\deg(L) = \tilde{C} \cdot b^*K_Y = C \cdot K_Y = 2\gamma - 2 - C^2 + 2\delta \quad (37)$$

we can rewrite the previous inequality in the following form:

$$\begin{aligned} 28 &\geq 2[2(\gamma - 1) - C^2 + 2\delta - 2p_g + 2] + C^2 - 2\delta \\ &= 2[\deg(L) - 2p_g + 2] + C^2 - 2\delta \\ &\geq 2 \operatorname{Cliff}(L) + C^2 - 2\delta \quad (\text{by (36)}) \\ &\geq 2 \operatorname{Cliff}(\tilde{C}) + C^2 - 2\delta \geq \gamma - 2 + \tilde{C}^2 + 2\delta \quad (\text{by (36) again}) \end{aligned}$$

Now, since $h^0(\mathcal{O}_{\tilde{C}}(\tilde{C})) \geq 2$, we have $\tilde{C}^2 \geq \frac{1}{2}\gamma + 1$, by Proposition 2.5(ii), because $\tilde{C}$ is a general curve. Therefore:

$$28 \geq \frac{3}{2}\gamma - 1 + 2\delta \geq \frac{3}{2}\gamma - 1$$

which is equivalent to $\gamma \leq 19$.

Let's prove (ii). After substitution of the inequality $K_Y^2 \geq 4\chi(\mathcal{O}_Y) - 16$ in (30) we obtain:

$$2\chi(\mathcal{O}_Y) + 32 - 2\delta \geq 4(\gamma - 1) - C^2$$

which, as before, can be replaced by the weaker:

$$34 \geq 4(\gamma - 1) - C^2 - 2p_g + 2\delta \quad (38)$$

Using again the expression (37) for the degree of L , we can write:

$$\begin{aligned} 36 &\geq 2(\gamma - 1) + \deg(L) - 2p_g + 2 \\ &\geq 2(\gamma - 1) + \operatorname{Cliff}(L) \\ &\geq 2(\gamma - 1) + \operatorname{Cliff}(\tilde{C}) \geq \frac{5}{2}(\gamma - 1) \end{aligned}$$

which is equivalent to $\gamma \leq 15$.

We now prove (iii). The hypothesis implies that $h^0(\omega_{\tilde{C}}L^{-1}) \geq 1$. We may assume that $h^0(\omega_{\tilde{C}}L^{-1}) = 1$ because otherwise we can proceed as in case (ii) and get $\gamma \leq 15$. Thus:

$$\gamma - \deg(\omega_{\tilde{C}}L^{-1}) = \gamma - C^2 + 2\delta = h^0(\tilde{C}, L) \geq p_g \quad (39)$$

where the inequality is true by the assumption $|K_Y - C| = \emptyset$. Therefore we can now write (38) as:

$$34 \geq 3(\gamma - 1) + h^0(\tilde{C}, L) - 1 - 2p_g \geq 3(\gamma - 1) - p_g - 1 \geq 2\gamma - 4$$

(we have used $p_g \leq \gamma$, which follows from (39)). This proves $\gamma \leq 19$. $\square$

The following is a notable special case of the theorem:

Corollary 6.2 *Let Y be a minimal surface of general type such that the canonical map is birational onto its image. Suppose that $C \subset Y$ is a nodal irreducible curve of geometric genus γ whose normalization has general moduli and such that $\dim(|C(-2p_1 - \dots - 2p_\delta)|) \geq 2$, where $p_1, \dots, p_\delta$ are the nodes of C . Then $\gamma \leq 19$.*

Proof. Y satisfies $K_Y^2 \geq 3\chi(\mathcal{O}_Y) - 10$ (Castelnuovo inequality, see [8]). Then we are in case (i) of Theorem 6.1. $\square$

Remark 6.3 The surface $Y \subset \mathbb{P}^6$ considered in Remark 3.4(ii) is minimal and satisfies

$$K_Y^2 = 16 = 4\chi(\mathcal{O}_Y) - 16$$

Therefore it shows that (ii) of Theorem 6.1 is sharp. It is unknown whether the bound on the genus given by 6.1(i) and 6.1(iii) is sharp.

The proof of Theorem 6.1 seems to suggest that the bound $\gamma \leq 19$ could be violated by a general curve of geometric genus γ moving in a non-trivial linear system on a minimal surface Y such that $K_Y^2 < 3\chi(\mathcal{O}_Y) - 10$. We will show that, under certain technical assumptions, this is actually not possible. The point is that a surface of this type is a double cover of a ruled surface and from this fact one can easily reduce to Theorems 4.1 and 4.2.

We collect in the following statement a few known facts that we will need.

Proposition 6.4 *Let Y be a minimal surface of general type. Denote by p_g its geometric genus. Assume that $K_Y^2 < 3\chi(\mathcal{O}_Y) - 10$. Then:*

- (i) $p_g \geq 4$.
- (ii) *There is a commutative diagram:*

$$\begin{array}{ccccc} Z & \xrightarrow{b} & Y & & \\ \downarrow \pi & & \downarrow & \searrow \varphi_K & \\ R & \longrightarrow & \Sigma & \hookrightarrow & \mathbb{P}^{p_g-1} \end{array} \quad (40)$$

where φ_K is the canonical rational map, which is generically finite and of degree 2; b is a blow-up, π is a degree 2 finite map, R is a birationally ruled surface, not necessarily minimal, and Σ is a surface of degree strictly smaller than $2(p_g - 1)$.

Proof. (i) The inequality is an obvious consequence of the assumption.

(ii) The existence of diagram (40) follows from Th. 5.5 of [8] and from Proposition 3.2 of [23]. $\square$

Theorem 6.5 *Let Y be a minimal surface of general type such that*

$$K_Y^2 < 3\chi(\mathcal{O}_Y) - 10$$

Let $C \subset Y$ be an irreducible curve of geometric genus $\gamma \geq 3$ having nodes $p_1, \dots, p_\delta$ and no other singularities. Assume that

- (i) *C is contained in a linear pencil Λ whose general member is an irreducible curve with nodes at $p_1, \dots, p_\delta$ and no other singularities.*
- (ii) *The normalization $\tilde{C}$ of C is a general nonsingular curve of genus γ .*

Let $C' \subset Z$ be the proper transform of C under the blow-up b whose existence is stated in Proposition 6.4(ii).

Again referring to diagram (40), assume further that:

- (iii) *there is a birational contraction $m : R \rightarrow F$ to a minimal model F of R such that $m(\pi(C'))$ is a nodal curve.*

Then Y is regular (i.e. it has irregularity $q = 0$) and $\gamma \leq 14$.

Proof. $\pi|_{C'} : C' \rightarrow \pi(C')$ is an isomorphism, because otherwise $\tilde{C}$ would be hyperelliptic or $\pi|_{C'}$ would define an irrational involution, and both possibilities contradict the fact that $\tilde{C}$ has general moduli (Proposition 2.5). The pencil Λ induces a pencil of curves in R containing $\pi(C')$, which in turn defines a pencil of curves in F containing $m(\pi(C'))$ as a member.

Assume that Y is irregular. In this case R is a non-rational birationally ruled surface. From Proposition 2.8 it follows that the pencil of nodal curves in R containing $\pi(C')$ defines an isotrivial fibration. This implies that the pencil Λ in Z defines an isotrivial fibration as well, contradicting Corollary 2.7. Therefore we have a contradiction, and it follows that Y is regular, and that R is rational.

Since $m(\pi(C'))$ is a nodal curve of geometric genus γ whose normalization is $\tilde{C}$, a general curve of genus γ , we are in the hypothesis of either Theorem 4.1, if $F = \mathbb{P}^2$, or of Theorem 4.2, if $F = F_n$. We conclude that $\gamma \leq 14$. $\square$

References

- [1] Arakelov S. Ju.: Families of algebraic curves with fixed degeneracy, *Math. USSR Izvestija* 5(1971), 1277-1302.
- [2] Araujo C.: Rationally connected varieties, *Contemporary Math.* 388 (2006), 1-16.
- [3] Araujo C., Kollar J.: Rational curves on varieties, in *Higher dimensional varieties and rational points (Budapest 2001)*, 13-68, Springer 2003.
- [4] Arbarello E.: Alcune osservazioni sui moduli delle curve appartenenti ad una data superficie algebrica, *Rend. Acc. Naz. Lincei* (VIII) 59 (1975), 725-732.
- [5] Arbarello E., Cornalba M.: Su una congettura di Petri, *Comment. Math. Helvetici* 56 (1981), 1-38.
- [6] Arbarello E., Sernesi E.: The equation of a plane curve, *Duke Math. J.* 46 (1979), 469-485.

- [7] Barth W., Peters C., Van de Ven A.: *Compact Complex Surfaces*, Springer Ergebnisse b. 4 (1984).
- [8] Beauville A.: L'application canonique pour les surfaces de type general, *Inventiones Math.* 55 (1979), 121-140.
- [9] Bruno A., Verra A.: M_{15} is rationally connected, in *Projective varieties with unexpected properties*, 51-65, Walter de Gruyter 2005.
- [10] Coppens M., Martens G.: Secant spaces and Clifford's theorem, *Compositio Math.* 78 (1991), 193-212.
- [11] Eisenbud D., Harris J.: The Kodaira dimension of the moduli space of curves of genus ≥ 23 , *Inventiones Math.* 90 (1987), 359-387.
- [12] Farkas G.: The geometry of the moduli space of curves of genus 23, *Math. Annalen* 318 (2000), 43-65.
- [13] Farkas G.: $\overline{\mathcal{M}}_{22}$ is of general type, *preprint* 2006.
- [14] Flamini F., Knutsen A., Pacienza G., Sernesi E.: Nodal curves with general moduli on K3 surfaces, *preprint*.
- [15] Griffiths Ph. and Harris J.: On the variety of special linear systems on a general algebraic curve, *Duke Math. J.* 47 (1980), 233-272.
- [16] Harris J. and Mumford D.: On the Kodaira dimension of the moduli space of curves, *Inventiones Math.* 67 (1982), 23-86.
- [17] Hartshorne R.: *Algebraic Geometry*, Springer Graduate Texts in Mathematics, v. 52 (1977).
- [18] Kleiman S.: Relative duality for quasi-coherent sheaves, *Compositio Math.* 41 (1980), 39-60.
- [19] Lange H.: Universal families of extensions, *J. of Algebra* 83 (1983), 101-112.
- [20] Mayer A.: Families of K3 surfaces, *Nagoya Math. J.* 48 (1972), 1-17.
- [21] Mori S., Mukai S.: The uniruledness of the moduli space of curves of genus 11, *Springer Lecture Notes in Math.* 1016 (1983), 334-353.

- [22] Mukai S.: Curves, $K3$ surfaces and Fano 3-folds of genus ≤ 10 . *Algebraic geometry and commutative algebra*, Vol. I, 357–377, Kinokuniya, Tokyo, 1988.
- [23] Persson U.: Double coverings and surfaces of general type, *Springer Lecture Notes in Math.* 687 (1978), 168-195.
- [24] Sernesi E.: *Deformations of Algebraic Schemes*, Springer Grundlehren b. 334 (2006).
- [25] Serrano F.: Fibred surfaces and moduli, *Duke Math. J.* 67 (1992), 407-421.
- [26] Serrano F.: Isotrivial fibred surfaces, *Annali di Mat. P. A.* (4) 171 (1996), 63-81.
- [27] Szpiro L.: Propriétés numériques du faisceau dualisant relatif, *Asterisque* 86 (1981), 44-78.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ ROMA TRE
 LARGO S.L. MURIALDO 1, 00146 ROMA
 sernesi@mat.uniroma3.it